

NEW VARIABLE STARS IN THE GLOBULAR CLUSTER NGC 6584¹

JOSEPH M. TODDY²

Department of Physics and Astronomy, University of Georgia, Athens, GA 30601

AND

ELLIOTT W. JOHNSON, ANDREW N. DARRAGH, AND BRIAN W. MURPHY

Department of Physics and Astronomy, Butler University, Indianapolis, IN, 46208 & SARA

ABSTRACT

Using the Southeastern Association for Research in Astronomy 0.6 meter telescope located at Cerro Tololo, we searched for variable stars in the southern globular cluster NGC 6584. We obtained images during 8 nights between 28 May and 6 July of 2011. After processing the images, we used the image subtraction package ISIS developed by Alard (2000) to search for the variable stars. We identified a total of 69 variable stars in our 10×10 arcmin² field, including 43 variables cataloged by Millis & Liller (1980) and 26 hereto unknown variables. In total, we classified 46 of the variables as type RRab, with a mean period of 0.56776 days, 15 as type RRc with a mean period of 0.30886 days, perhaps one lower amplitude type RRe, with a period of 0.26482 days, 4 eclipsing binaries, and 3 long period ($P > 2$ days) variable stars. As many as 15 of the RRab Lyrae stars exhibited the Blažhko Effect. Furthermore, the mean periods of the RR Lyrae types, the exhibited period/amplitude relationship, and the ratio of $N_c/(N_{ab} + N_c)$ of 0.25 are consistent with an Oosterhoff Type I cluster. Here we present refined periods, V-band light curves, and classifications for each of the 69 variables, as well as a color-magnitude diagram of the cluster.

Subject headings: stars: variables: general–Galaxy: globular clusters: individual: NGC 6584

1. INTRODUCTION

Globular clusters provide an ideal controlled environment to study variable stars and hence stellar evolution since the stars in the cluster presumably formed out of the same cloud of gas and dust, have the same composition, have the same distance, and have the same age (Smith 2004). In addition to understanding the important aspects of late-stage stellar evolution in a controlled environment, variable stars in globular clusters are one of the primary standard candles that are used to measure the distances to globular clusters in the Milky Way and nearby galaxies. They also provide a method to easily observe thousands of stars at once, a useful characteristic when searching for variable stars.

NGC 6584 is a relatively understudied globular cluster located 13.5 Kpc from the Sun in the southern constellation Telescopium (Harris 1996). It is an Oosterhoff type I cluster (Oosterhoff 1939) and Sarajedini & Forrester (1995) have estimated its metallicity at $[\text{Fe}/\text{H}] = -1.44 \pm 0.16$. Early studies of this cluster were performed by Bailey (1924) who identified nine potential variable stars in its vicinity. Millis & Liller (1980) published the locations of 48 variable stars in NGC 6584. They utilized photographic plates in their observations and analyses. As a result, the light curves that they present are relatively incomplete and the periods that they provide are somewhat imprecise. Sarajedini & Forrester (1995) confirmed these 42 RR Lyrae variables and tentatively

identified 56 possible yet undiscovered RR Lyraes by analyzing the variation in the frame-to-frame dispersion in V and B-V as a function of V for the stars in the instability strip of their color-magnitude diagram (CMD). Those stars which showed a large range in V dispersion were identified as possible RR Lyraes. In the same year, Samus et al. (1995), also using photographic photometry, again confirmed Millis & Liller's 48 variables. Since the results of Sarajedini & Forrester (1995) and Samus et al. (1995) image subtraction software and better CCDs have become available.

With the combination of CCD images, an observation window spanning more than 5 weeks, and an image subtraction method developed by Alard (2000), we are able to detect and analyze hereto undiscovered variable stars and produce unprecedented high-quality light curves and periods for each variable. In this study we use Alard's image subtraction method to search the central 10×10 arcmin² of globular cluster NGC 6485 for variable stars from observations obtained in May, June, and July of 2011. We present a preliminary color-magnitude diagram for the cluster, as well as complete light curves and periods for 43 of Millis & Liller's variables and for 26 newly identified variables.

2. OBSERVATIONS AND REDUCTION

We obtained images using the Southeastern Association for Research in Astronomy (SARA) 0.6 meter telescope at Cerro Tololo Interamerican Observatory (CTIO). Observations were made on the nights of 28 May; 2, 22, 25, and 28 June; 2, 5, and 6 July 2011 using an Apogee Alta E6 camera with a 1024×1024 pixel Kodak KAF1001E chip, a gain of 1.5 electron per ADU count, and an RMS noise of 8.9 electrons. We used 1×1 binning resulting in an image scale of $0.61''/\text{px}$ resulting

¹ Based on observations obtained with the SARA Observatory 0.6 m telescope at Cerro Tololo, which is owned and operated by the Southeastern Association for Research in Astronomy (<http://www.saraobservatory.org>).

² Southeastern Association for Research in Astronomy (SARA) NSF-REU Summer Intern
 Electronic address: bmurphy@butler.edu




FIG. 1.— The inner 5×5 arcmin² of the **ref.fits** image used for all nights of the analysis/image subtraction. This image is a combination of the best seeing images, with the combined seeing near $1.3''$. Note that the 6 brightest stars in the image are overexposed.

in a 10×10 arcmin² field of view. The CCD was kept at a constant temperature of -25 Celsius. The typical seeing was $1.7''$ and ranged from 1.2 to $2.8''$. On all nights of observation, 240 second exposures were taken using a Bessel V filter. The exposure time allowed us to detect dimmer variables but caused 6 of the stars in the field many to be overexposed. Most of these overexposed stars appeared to be in the foreground of the cluster. On the nights of 2, 5, and 6 July 2011, additional exposures were taken using Bessel B, R, and I filters with exposure times of 300, 90, and 50 seconds respectively. All images were debiased, flat-fielded, dark subtracted, and cleaned of hot, cold, and bad pixels using MaximDL.

3. ANALYSIS

Image Subtraction was accomplished using the ISIS-2.2 package (Allard & Lupton 1998; Allard 2000). ISIS accounts for changes in seeing conditions by convolving a high quality reference image to the point spread functions of each individual frame and then subtracting the two images to determine any change in intensity. The package contains seven c-shell scripts and three parameter files that were used in our analysis. The different c-shell scripts are **dates**, **interp** (registration and interpolation), **ref** (build reference frame), **subtract** (subtracts images from reference), **detect** (stacks subtracted images), **find** (finds variables above a user-defined threshold), and **phot** (produces light curves for all detected possible variables), along with three parameter files.

We began by running a modified version of the **dates** script, which identified and registered each image file in the image directory. We then ran the **interp** script which aligned each image to a previously chosen alignment image. Next, the **ref** script combined five of our highest quality images into a reference frame (Fig. 1). In order to maintain consistent and comparable values for relative flux across nights, one reference image was




FIG. 2.— The inner 5×5 arcmin² of the **var.fits** frame output from ISIS for all eight nights from SARA CTIO. The relative amount of variation is indicated by the brightness of the star. The 6 overexposed stars show significant variation only because ISIS is unable to fit an accurate point spread function to them.

created using five of the best images (those with seeing between 1.2 and $1.3''$). We then ran the **subtract** script, which first convolves the high-quality reference frame to the point-spread function of the image to be subtracted (in order to compensate for variations in seeing over the course of the observations), and then subtracts the convolved reference frame from each image. These subtracted images were then stacked using the **detect** script creating a **var.fits** image which shows the degree of variability across the observation run for each star in the cluster as shown in Fig. 2. We then used the **find** script to identify variable stars above a certain user-defined threshold called **SIGTHRESH**. We used a **SIGTHRESH** of 0.03 in our analysis to ensure that all possible variables were found. This value typically resulted in 1500 possible variables. Finally, the **phot** script was used to generate light curve data files for each identified variable. These light curves allowed us to eliminate false positives from the **find** script. Variable stars produce very distinct light curves, and thus false positives due to noise and variations in seeing were easily identified amongst these plots and discarded.

Periods were determined, when possible, based on combined photometric data from all eight nights. We used the period finding software **AVE** (Análisis de Variabilidad Estelar, Analysis of Estellar Variability) from Grup d'Estudis Astronòmics. **AVE** uses the Lomb-Scargle algorithm (Scargle 1982) to determine the most probable period for a given set of data. Due to the more than 5 week observation span, we were able to determine the periods of most of the variables with periods less than 1 day to an accuracy of 10^{-5} days. Those variables exhibiting the Blazhko Effect were less accurate due to modulation of their light curves.

Millis & Liller's (1980) variables were identified visually and astrometrically and then confirmed by compar-

TABLE 1
CLASSIFICATION SUMMARY

Variable Type	Count	Period (days)
RRab	46	0.56776
RRc	15	0.30886
RRe	1	0.26482
Eclipsing	4	0.32431
LPV	3	> 20

ing their periods with our own. Five of the variables identified by Millis & Liller were outside the field of view of our images thus were not be observed.

4. RESULTS

Table 1 summarizes our findings by classification. Table 2 list the positions, periods, and classification of all of the variables that we identified. In Table 2 variables previously identified by Millis & Liller (1980) are labeled with a V and the number they assigned in the first column. Newly discovered variables are labeled as NV and are in order of increasing right ascension. We detected all the Millis & Liller (1980) variables that were within our 10×10 arcmin² field. Their variables 24, 27, 40, 43, and 44 were outside our field of view. We did note that variable 45 was misidentified on the finder chart in Millis & Liller (1980), thus the position we list for of this variable has been shifted by several arc-seconds in Table 2. We found periods for variables V12, V19, and 39 that Millis & Liller were unable to determine. We also were able to eliminate those periods where multiple possible periods were listed; specifically V5, V20 and V35. Besides the increased accuracy of our periods the only significant change from Millis and Liller was V13 which we found to be 0.34549 days as compared to 0.39 days.

Each detected variable was classified based on the shape of its light curve, its period, and its maximum amplitude. In addition to the Miller & Liller variables we found 26 additional variables in our field, most of which are RR Lyrae variables, long period variables, or eclipsing variables. Of the total 69 variables detected, 46 were classified as RRab stars, 15 as RRc, 1 possible RRe, 4 eclipsing binaries, and 3 long period variables. From the fraction $N_c/(N_{ab} + N_c)$ we confirm that the cluster is of Oosterhoff Type I. RRab stars are characterized by large amplitudes and periods between 0.4 and 0.8 days. RRc stars have smaller amplitudes and periods between 0.2 and 0.4. NV9 has the characteristics of a type RRe variable (Alcock et al. 1996; Bono et al. 1997). It has a lower amplitude and period than type RRc stars. Eclipsing binary stars are easily identified by their unique light curves. As many as 15 of the RRab stars appeared to exhibit the so-called Blažhko effect (Blažhko 1907; Smith 2004), which is characterized by a long term modulation in amplitude and period. These variables are noted with an asterisk in Table 2. Their periods are likely to be less accurate given their behaviour. We have not assigned periods to the long period variables at the tip of the asymptotic giant branch because further observations will be needed to establish the their periods. We assign 'LPV' to these variables with a period greater than 2 days.

TABLE 2
VARIABLES STARS IN NGC 6584

V#	RA (h,m,s)	Dec (°,'")	Period (d)	Type
V1	18 18 28.82	-52 13 27.0	0.49520	RRab
V2	18 18 42.20	-52 12 30.0	0.54465	RRab
^a V3	18 18 42.14	-52 13 16.2	0.47162	RRab
V4	18 18 40.30	-52 13 25.5	0.54377	RRab
^b V5	18 18 40.49	-52 13 29.1	0.29725	RRc
^a V6	18 18 34.45	-52 12 7.8	0.62619	RRab
V7	18 19 1.34	-52 11 55.3	0.54748	RRab
V8	18 18 55.62	-52 13 35.3	0.30511	RRc
^a V9	18 18 43.55	-52 12 2.7	0.49513	RRab
V10	18 18 37.07	-52 11 36.7	0.69432	RRab
^a V11	18 18 35.42	-52 13 10.8	0.53945	RRab
^a V12	18 18 35.84	-52 12 32.6	0.59505	RRab
V13	18 18 35.85	-52 14 32.3	0.34549	RRc
^a V14	18 18 41.20	-52 13 43.2	0.59954	RRab
V15	18 18 23.56	-52 12 41.9	0.63632	RRab
V16	18 18 42.04	-52 12 58.6	0.58220	RRab
V17	18 12 41.87	-52 13 22.3	0.56338	RRab
^a V18	18 18 32.99	-52 13 18.3	0.55789	RRab
^a V19	18 18 40.66	-52 13 6.4	0.54223	RRab
^{a,b} V20	18 18 28.15	-52 12 57.9	0.52530	RRab
V21	18 18 34.48	-52 13 55.1	0.54025	RRab
V22	18 18 38.02	-52 12 50.5	0.55559	RRab
V23	18 18 30.76	-52 13 10.0	0.59812	RRab
V25	18 18 56.91	-52 11 12.8	0.45336	RRab
V26	18 18 33.62	-52 13 52.1	0.56970	RRab
V28	18 18 48.09	-52 12 19.8	0.49815	RRab
V29	18 18 41.49	-52 11 40.5	0.60720	RRab
^a V30	18 18 42.85	-52 9 31.9	0.51375	RRab
V31	18 18 30.61	-52 13 38.5	0.56831	RRab
V32	18 18 40.71	-52 14 5.4	0.59515	RRab
V33	18 18 26.81	-52 11 55.2	0.28595	RRc
V34	18 18 30.11	-52 12 16.4	0.78010	RRab
^b V35	18 18 36.82	-52 12 48.4	0.55550	RRab
V36	18 18 38.94	-52 14 0.2	0.52454	RRab
V37	18 18 26.06	-52 14 28.1	0.29786	RRc
^a V38	18 18 26.81	-52 11 55.1	0.55159	RRab
V39	18 18 19.48	-52 10 47.2	0.33141	EB
^a V41	18 18 40.70	-52 13 4.9	0.63992	RRab
V42	18 18 39.88	-52 11 55.3	0.57675	RRab
^c V45	18 18 28.75	-52 12 30.8	0.50823	RRab
V46	18 18 25.08	-52 12 35.3	0.52555	RRab
V47	18 18 38.32	-52 13 20.8	0.27666	RRc
V48	18 18 38.55	-52 13 26.1	0.30706	RRc
NV1	18 18 6.83	-52 12 25.0	0.25531	EB
NV2	18 18 19.26	-52 12 38.3	0.29297	RRc
NV3	18 18 27.84	-52 12 59.4	0.75210	RRab
NV4	18 18 27.99	-52 12 55.7	0.52384	RRab
NV5	18 18 28.19	-52 14 7.0	0.34724	RRc
NV6	18 18 32.16	-52 14 43.2	0.64406	RRab
NV7	18 18 32.79	-52 12 24.7	0.56583	RRab
NV8	18 18 33.14	-52 10 15.5	0.30807	RRc
NV9	18 18 34.47	-52 12 49.1	0.26482	RRe
NV10	18 18 35.57	-52 13 31.4	0.63930	RRab
NV11	18 18 35.66	-52 12 20.6	> 20	LPV
NV12	18 18 35.67	-52 13 7.1	0.58387	RRab
NV13	18 18 37.69	-52 12 59.8	> 20	LPV
NV14	18 18 37.79	-52 13 6.2	> 20	LPV
^a NV15	18 18 38.16	-52 12 54.8	0.53741	RRab
NV16	18 18 38.42	-52 12 56.0	0.63630	RRab
NV17	18 18 38.58	-52 12 21.4	0.49644	RRab
NV18	18 18 38.77	-52 13 14.1	0.34180	RRc
^a NV19	18 18 39.48	-52 14 32.8	0.58572	RRab
NV20	18 18 39.93	-52 11 21.3	0.31722	RRc
NV21	18 18 40.01	-52 13 9.1	0.30911	RRc
NV22	18 18 40.81	-52 10 32.5	0.33625	RRc
^a NV23	18 18 41.98	-52 12 53.7	0.55739	RRab
NV24	18 18 42.72	-52 12 28.9	0.27374	RRc
NV25	18 18 44.98	-52 8 10.7	0.24689	EB
NV26	18 18 51.42	-52 13 30.8	0.46364	EB

^a variable exhibiting Blažhko Effect.^b variables with significantly revised periods.^c misidentified by Millis & Liller (1980).




FIG. 3.— Phased light curves for identified variables with periods under a day. The vast majority of these variables are RR Lyrae stars.




FIG. 3.— (Continued) Phased light curves for identified variables with periods under a day. The vast majority of these variables are RR Lyrae stars.




FIG. 3.— (Continued) Phased light curves for identified variables with periods under a day. The vast majority of these variables are RR Lyrae stars.




FIG. 3.— (Continued) Phased light curves for identified variables with periods under a day. The vast majority of these variables are RR Lyrae stars.




FIG. 3.— (Continued) Phased light curves for identified variables with periods under a day. The vast majority of these variables are RR Lyrae stars.




FIG. 4.— Light curves for the 3 long period variables found in our analysis.

Cluster membership was determined using a color magnitude diagram (CMD). The CMD was constructed from 137 V and 155 R frames from several nights to determine magnitudes, colors, and cluster membership of the variables. DAOPHOT was used to determine V and R magnitudes of the stars (Stetson 1987). In the CMD shown in Figure 5 RRab and RRC variables are indicated by the red and blue colored points, respectively. RRC variables tend to lie on the warmer (left) side of the instability strip whereas RRab variables are to the cooler (right) of the instability strip. Scatter in the variables is partly due to blending of the variables with other stars, our averaging of several hundred images before using DAOPHOT, and possible variable reddening in the direction of the cluster.

Six of our variables, V11, V22, NV7, NV19, NV20, and NV21 were blended with other stars and show significant offset to the right of the instability strip and are likely cluster members. Four of the RRab variables V15, V41, NV4, and NV15 were significantly brighter than the RR Lyrae variables found in the instability strip of the cluster and are suspected not to be cluster members. From the

CMD we are able to determine that 2 of the eclipsing variables to the right of the giant branch are not cluster members. The 3 long period variables that we found all lie near the tip of the asymptotic giant branch. The remainder of the RR Lyrae stars lie very near or in the instability strip and thus are likely cluster members.

5. CONCLUSIONS

Over a 39 day time span from late May to early July of 2011 we observed the globular cluster NGC 6584 for 8 nights using the SARA 0.6 meter telescope at CTIO. Using the image subtraction software ISIS we were able to find precise periods of previously known variables and 26 newly discovered variables stars, most of which are RR Lyrae stars. Within our 10×10 arcmin² field of view we detected all the variables listed by Millis & Liller (1980) bringing the total to 74 variables in the vicinity of the cluster. Of all the RR Lyrae variables all but 4 appeared to be members of the cluster. Two of the 4 eclipsing binaries are likely to be cluster members. In total, we classified 46 of the variables as type RRab, with a mean period of 0.56776 days, 15 as type RRC with a mean




FIG. 5.— Color-magnitude diagram of NGC 6584, with variable stars indicated. Note that the RRab's (red) variables are bluer than the RRc's (blue). All but 6 of the variables found appear to be cluster members. The two eclipsing variables to the right of the giant branch of the cluster are likely not cluster members. The 2 RRc and 4 RRab Lyrae variables just to the right of the instability strip are all blends. The 4 brightest RRab's may be foreground objects.

period of 0.30886 days, perhaps one lower amplitude type RRe, with a period of 0.26482 days, 4 eclipsing binaries, with a mean period of 0.3243 days, and 3 long period ($P > 2$ days) variable stars. As many as 15 of the RRab Lyrae stars exhibited the Blazhko Effect. The ratio of $N_c/(N_{ab} + N_c)$ of 0.25 is consistent with cluster being of type Oosterhoff Type I.

We thank C. Alard for making ISIS 2.2 publically avail-

able. This project was funded in part by the National Science Foundation Research Experiences for Undergraduates (REU) program through grant NSF AST-1004872. Additionally A. Darragh, E. Johnson, B. Murphy were partially funded by the Butler Institute for Research and Scholarship. The authors also thank F. Levinson for a generous gift enabling Butler University's membership in the SARA consortium.

REFERENCES

- Alard, C., & Lupton, R. H. 1998, *ApJ*, 503, 325
 Alard, C. 2000, *A&AS*, 144, 363
 Alcock, C., Allsman, R. A., Axelrod, T. S., et al. 1996, *AJ*, 111, 1146
 Bailey, S. I. 1924, *Harvard College Observatory Bulletin* 801, pp.7-8
 Blažko, S. 1907, *Astronomische Nachrichten*, 175, 325
 Bono, G., Caputo, F., Cassisi, S., Incerpi, R., & Marconi, M. 1997, *ApJ*, 483, 811
 Harris, W. E. 1996, *AJ*, 112, 1487
 Millis, A. J. & Liller, M. H. 1980, *AJ*, 85, 3
 Oosterhoff, P. T. 1939, *The Observatory*, 62, 104
 Samus, N. N. et al. 1995, *AJ*, 21,4
 Sarajedini, A. & Forrester, W. L. 1995, *AJ*, 109, 3
 Scargle, J. D. 1982, *ApJ*, 263, 835
 Smith, H. A. 2004, *RR Lyrae Stars*, by Horace A. Smith, ISBN 0521548179. Cambridge, UK: Cambridge University Press
 Stetson, P. B. 1987, *PASP*, 99, 191